



## Adult Tobacco Use in Pipestone County

### Findings from the 2015 Southwest Minnesota Healthy Communities Survey

The 2015 Southwest Minnesota Healthy Communities Survey was conducted to learn about the health and health habits – physical activity, nutrition, tobacco use, dental health, and mental health – of residents living in 16 counties in southwest Minnesota. Southwest Health and Human Services contracted the Wilder Research to produce summary reports for each of the six counties in their service area. This fact sheet summarizes survey results related to the tobacco use of adult residents in Pipestone County.


The survey was conducted April through July 2015 and was sponsored in part by the participating counties' grants from the Minnesota Department of Health's Statewide Health Improvement Program (SHIP).

In the following report, when it is stated that respondents' answers to a survey item increased or decreased, this refers to a change of 10 percentage points or more in respondents' answers from the 2010 to the 2015 county survey. Similarly, when response percentages are reported as higher or lower in the county compared to the 16-county region this refers to a difference of 10 percentage points or more.

### *Current tobacco use*

Sixty-one percent of Pipestone County residents have never smoked, and 24 percent are former smokers. Fifteen percent of Pipestone County respondents are current smokers.

### RESPONDENTS' SMOKING STATUS



### *E-cigarettes*


Two percent of respondents said they are e-cigarette users. About half (46%) of respondents reported ever having seen someone smoke an e-cigarette in Minnesota. Respondents were most likely to have seen someone use an e-cigarette in a restaurant or bar, or in someone's home or car (including their own).

### *Quitting*

Of the 39 percent who are current and former smokers, 43 percent said they stopped smoking for one day or longer in the past year because they were trying to quit.

Of former and current smokers who had stopped smoking for one day or longer during the past year because they were trying to quit, the most common quitting method used was to quit without any assistance from nicotine replacement medications, counseling, or other forms of quitting support.

**RESPONDENTS' USE OF VARIOUS TOBACCO CESSATION METHODS (OF CURRENT AND FORMER SMOKERS WHO HAD TRIED TO QUIT IN THE PAST 12 MONTHS)**



“Quitting without any assistance” was more commonly used among successful quitters (those who are now former smokers and who quit within the past 12 months) than among unsuccessful quitters (current smokers who tried to quit at least once during the past 12 months).


Residents of Pipestone County were less likely in 2015 than in 2010 to have stopped smoking for one day or longer because they were trying to quit.

**Exposure to smoking**

Six percent of Pipestone County respondents allow regular smoking in their homes (by themselves or someone else) compared with 10 percent of respondents from the region. Twelve percent report they rode in a vehicle with someone who was smoking in the past week and 26 percent of respondents say they were exposed to secondhand smoke in a public (outdoor) place in Minnesota during the past week.

Since 2010, exposure to secondhand smoke in public spaces has decreased.

**RESPONDENTS' EXPOSURE TO SECONDHAND SMOKE IN THE PAST 7 DAYS**



**Compared with the region...**

There are no significant differences in tobacco use in Pipestone County compared with tobacco use in the 16-county region.



**Wilder Research**

Information. Insight. Impact.

AMHERST H  
WILDER  
FOUNDATION  
ESTABLISHED 1906

Here for good.

451 Lexington Parkway North  
Saint Paul, Minnesota 55104  
651-280-2700  
www.wilderresearch.org

**For more information**

This summary presents findings from the 2015 Southwest Minnesota Healthy Communities Survey. For more information about the survey, contact Nicole MartinRogers at Wilder Research, 651-280-2682. For access to other reports, visit <http://www.swmhhs.com/public-health-assessment-and-planning/>

Authors: Monzong Cha, Anna Bartholomay, and Nicole MartinRogers  
APRIL 2016